

Non-Abelian Anyons in the Quantum Hall Effect

Andrea Cappelli

(INFN and Physics Dept., Florence)


with L. Georgiev (Sofia), G. Zemba (Buenos Aires), G. Viola (Florence)

Outline

- Incompressible Hall fluids: bulk & edge excitations
- CFT description
- Non-Abelian statistics
- Partition function
- Signatures of non-Abelian statistics:
 - Coulomb blockade & thermopower

Quantum Hall Effect

- 2 dim electron gas at low temperature $T \sim 10$ mK
and high magnetic field $B \sim 10$ Tesla





- **Conductance tensor** $J_i = \sigma_{ij} E_j$, $\sigma_{ij} = R_{ij}^{-1}$, $i, j = x, y$
- **Plateaux:** $\sigma_{xx} = 0$, $R_{xx} = 0$ **no Ohmic conduction** \rightarrow gap
 $\sigma_{xy} = R_{xy}^{-1} = \frac{e^2}{h} \nu$, $\nu = 1(\pm 10^{-8}), 2, 3, \dots, \frac{1}{3}, \frac{2}{5}, \dots, \frac{5}{2}$,
- **High precision & universality**
- Uniform density ground state: $\rho_o = \frac{eB}{hc} \nu$

Incompressible fluid

Laughlin's quantum incompressible fluid



Electrons form a droplet of fluid:

- incompressible = gap
- fluid = $\rho(x, y) = \rho_o = \text{const.}$



$\mathcal{D}_A = BA/\Phi_o$, # degenerate orbitals = # quantum fluxes, $\Phi_o = \frac{hc}{e}$

filling fraction: $\nu = \frac{N}{\mathcal{D}_A} = 1, 2, \dots, \frac{1}{3}, \frac{1}{5}, \dots$ density for quantum mech.



Laughlin's wave function

$$\Psi_{gs}(z_1, z_2, \dots, z_N) = \prod_{i < j} (z_i - z_j)^{2k+1} e^{-\sum |z_i|^2 / 2} \quad \nu = \frac{1}{2k+1} = 1, \frac{1}{3}, \frac{1}{5}, \dots$$

- $\nu = 1$ filled Landau level: obvious gap $\omega = \frac{eB}{mc} \gg kT$
- $\nu = \frac{1}{3}$ non-perturbative gap due to Coulomb interaction

➔ effective theories

- quasi-hole = elementary vortex $\Psi_\eta = \prod_i (\eta - z_i) \Psi_{gs}$

➔ fractional charge $Q = \frac{e}{2k+1}$ & statistics $\frac{\theta}{\pi} = \frac{1}{2k+1}$



$$\Psi_{\eta_1, \eta_2} = (\eta_1 - \eta_2)^{\frac{1}{2k+1}} \prod_i (\eta_1 - z_i) (\eta_2 - z_i) \Psi_{gs}$$

Anyons

vortices with long-range topological correlations


Conformal field theory of edge excitations

The edge of the droplet can fluctuate: edge waves are massless



- edge \sim Fermi surface: linearize energy $\varepsilon(k) = \frac{v}{R}(k - k_F)$, $k = 0, 1, \dots$
 - relativistic field theory in 1+1 dimensions, chiral (X.G.Wen '89)
- ➔ chiral compactified c=1 CFT (chiral Luttinger liquid)

CFT descriptions of QHE



- same function by analytic continuation from the circle:
 - both equivalent to Chern-Simons theory in 2+1 dim (Witten '89)
- $c=1$ Luttinger CFT:
 - wavefunctions: spectrum of anyons and braiding
 - edge correlators: physics of conduction experiments

Non-Abelian fractional statistics

- $\nu = \frac{5}{2}$ described by Moore-Read "Pfaffian state" ~ Ising CFT x boson
- Ising fields: I identity, ψ Majorana = electron, σ spin = anyon
- fusion rules:

- $\psi \cdot \psi = I$ 2 electrons fuse into a Bosonic bound state


- $\sigma \cdot \sigma = I + \psi$ 2 channels of fusion = 2 conformal blocks

$$\langle \sigma(0)\sigma(z)\sigma(1)\sigma(\infty) \rangle = a_1 F_1(z) + a_2 F_2(z) \quad \text{Hypergeometric}$$


➔ state of 4 anyons is two-fold degenerate (Moore, Read '91)

- statistics of anyons ~ analytic continuation ➔ 2x2 matrix

$$\begin{pmatrix} F_1 \\ F_2 \end{pmatrix} (ze^{i2\pi}) = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} F_1 \\ F_2 \end{pmatrix} (z)$$



$$\begin{pmatrix} F_1 \\ F_2 \end{pmatrix} ((z-1)e^{i2\pi}) = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} F_1 \\ F_2 \end{pmatrix} (z)$$



(all CFT redone for Q. Computation: M. Freedman, Kitaev, Nayak, Slingerland, ..., 00'-10')

Topological quantum computation

- qubit = two-state system $|\chi\rangle = \alpha|0\rangle + \beta|1\rangle$
- QC: perform $U(2^n)$ unitary transformations in n qubit Hilbert space
- Proposal: (Kitaev; M. Freedman; Nayak; Simon; Das Sarma '06)

use non-Abelian anyons for qubits and operate by braiding

4-spin system $\alpha|F_1\rangle + \beta|F_2\rangle$ is 1 qubit (2n-spin has dim 2^{n-1})

- anyons topologically protected from decoherence (local perturbations)
- more stable but more difficult to create and manipulate
 - ➔ great opportunity
 - ➔ new experiments and model building

Models of non-Abelian statistics

- Study Rational CFTs with non-Abelian excitations:

- best candidate: Pfaffian & its generalization, the Read-Rezayi states

$$\nu = 2 + \frac{k}{k+2}, \quad \left\{ \begin{array}{l} k = 2, 3, \dots \\ M = 1 \end{array} \right. \quad U(1)_{k+2} \times \frac{SU(2)_k}{U(1)_{2k}}$$

- alternatives: other (cosets of) non-Abelian affine groups $U(1) \times \frac{G}{H}$

- Identify their N sectors of fractional charge and statistics

- Abelian (electron) & non-Abelian (quasi-particles)

- Compute physical quantities that could be signatures of non-Abelian statistics:


- Coulomb blockade conductance peaks
- thermopower & entropy



use partition function

- quantity defining Rational CFT (Cardy '86; many people)
- complete inventory of states (bulk & edge)
- modular invariance as building principle:
 - S matrix and fusion rules
 - further modular conditions for charge spectrum
 - straightforward solution for any non-Abelian state $U(1) \times \frac{G}{H}$
 - useful to compute physical quantities
- Inputs:
 - non-Abelian RCFT (i.e. $\frac{G}{H}$)
 - Abelian field representing the electron "simple current"
- Output is unique

Annulus partition function



$$i2\pi\tau = -\beta\frac{v}{R} + it, \quad \beta = \frac{1}{k_B T}$$

$$i2\pi\zeta = \beta(-V_o + i\mu)$$

$$Z_{\text{annulus}} = \sum_{\lambda=1}^p |\theta_{\lambda}(\tau, \zeta)|^2, \quad \theta_{\lambda}(\tau, \zeta) = \text{Tr}_{\mathcal{H}(\lambda)} \left[e^{i2\pi\tau(L_0 - c/24) + i2\pi\zeta Q} \right]$$

modular invariance conditions

geometrical properties & physical interpretation

(A. C., Zemba, '97)

$$T^2 : Z(\tau + 2, \zeta) = Z(\tau, \zeta), \quad L_0 - \bar{L}_0 = \frac{n}{2} \quad \text{half-integer spin excitations globally}$$

$$S : Z\left(\frac{-1}{\tau}, \frac{-\zeta}{\tau}\right) = Z(\tau, \zeta), \quad \text{completeness} \quad \theta_{\lambda}\left(\frac{-1}{\tau}\right) = \sum_{\lambda'} S_{\lambda\lambda'} \theta_{\lambda'}(\tau) \quad \text{S matrix}$$


$$U : Z(\tau, \zeta + 1) = Z(\tau, \zeta), \quad Q - \bar{Q} = n \quad \text{integer charge excitations globally}$$

$$V : Z(\tau, \zeta + \tau) = Z(\tau, \zeta), \quad \Delta Q = \nu \quad \text{add one flux: spectral flow}$$

$$\theta_{\lambda}(\zeta + \tau) \sim \theta_{\lambda+1}(\tau)$$

Disk partition function

Annulus \rightarrow Disk (w. bulk q -hole $\bar{Q} = \frac{\lambda}{p}$)



$$Z_{\text{annulus}} \rightarrow Z_{\text{disk}}, \lambda = \theta_{\lambda}(\tau, \zeta)$$

$$\theta_{\lambda}(\tau, \zeta) = K_{\lambda}(\tau, \zeta; p) = \frac{1}{\eta} \sum_n e^{i2\pi \left[\tau \frac{(np+\lambda)^2}{2p} + \zeta \frac{np+\lambda}{p} \right]}, \quad \nu = \frac{1}{p}, \quad c = 1$$

- U : $Q - \bar{Q} = n$ sectors with charge $Q = \frac{\lambda}{p} + n$

basic quasiparticle + n electrons

- T^2 : electrons have half-integer dimension (=J),
and integer relative statistics with all excitations
- # sectors $p = \dim(S_{\lambda\lambda'}) = \text{Wen's topological order}$

\rightarrow we recover phenomenological conditions on the spectrum

Pfaffian & Read-Rezayi states

$$Z_{annulus}^{RR} = \sum_{\ell=0}^k \sum_{a=0}^{\hat{p}-1} |\theta_a^\ell(\tau, \zeta)|^2, \quad Z_{disk}^{RR} = \theta_a^\ell = \sum_{\beta=0}^{k-1} K_{a+\beta(k+2)} \chi_{a+2\beta}^\ell$$

Ex: Pfaffian (k=2)

ground state + electrons

$$Z_{annulus}^{Pfaffian} = |K_0 I + K_4 \psi|^2 + |K_0 \psi + K_4 I|^2 + |(K_1 + K_{-3}) \sigma|^2 \\ + |K_2 I + K_{-2} \psi|^2 + |K_2 \psi + K_{-2} I|^2 + |(K_3 + K_{-1}) \sigma|^2$$

non-Abelian quasiparticle

- K_λ charge parts $Q = \frac{\lambda}{4} + 2n$
- I, ψ, σ Ising parts (Majorana fermion)
 - 6 sectors
 - also $Q = 0, \pm \frac{1}{2}$ Abelian excitations


- charge and neutral $q, \#$'s are coupled by "parity rule"
- but S-matrix of θ_a^ℓ is factorized $S_{al, a'l'} \sim e^{i2\pi a a' N/M} S_{ll'}$

- generalization to other N-A models: (A.C, G. Viola, '10)


- Non-Abelian Fluids $U(1) \times SU(2)_k$
- Anti-Read-Rezayi $U(1) \times \overline{SU(2)_k}$
- Bonderson-Slingerland $U(1) \times \text{Ising} \times SU(n)_1$
- N-A Spin Singlet state $U(1)_q \times U(1)_s \times \frac{SU(3)_k}{U(1)^2}$

➔ unique result once RCFT and electron field have been chosen

Experiments on non-Abelian statistics




(a)



(b)

- (a) interference of edge waves (Chamon et al. '97; Kitaev et al 06)
 - ➔ Aharonov-Bohm phase, checks fractional statistics
 - experiment is hard (Goldman et al. '05; Willett et al '09)
- (b) electron tunneling into the droplet (Stern, Halperin '06)
 - ➔ Coulomb blockade conductance peaks (Ilan, Grosfeld, Schoutens, Stern '08)
 - check quasi-particle sectors (Stern et al.; A.C. et al. '09 - '10)
- ➔ Thermopower (Cooper, Stern; Yang, Halperin '09; Chickering et al. '10)

Thermopower



- fusion of n non-Abelian quasiparticles:

- multiplicity $\sim (d_\ell)^n$, $n \rightarrow \infty$

- Entropy: $S(T = 0) \sim n \log(d_\ell)$, $d_\ell = \frac{s_{\ell 0}}{s_{00}} > 1$ (quantum dimension)

from Z: $S = \left(1 - \tau \frac{d}{d\tau}\right) \log \left[\theta_a^\ell(\tau, \zeta) \overline{\theta_0^0(\tau, \zeta)} \right] \sim \log \frac{s_{\ell 0}}{s_{00}} + S_{\text{Edge}}, \quad \tau \sim \frac{\beta}{R} \rightarrow 0$

- put temperature ∇T and potential ∇V_o gradients between the edges

- at equilibrium: $J = -\sigma \nabla V_o - \alpha \nabla T = 0$

- thermopower

$$S_{\text{Seebeck}} = \frac{\alpha}{\sigma} = -\frac{\nabla V_o}{\nabla T} = \frac{S}{eN_e}$$

(Cooper, Stern;
Yang, Halperin '09)

- it could be observable by varying B off the plateau center

$$S_{\text{Seebeck}} = \left| \frac{B - B_o}{e^* B_0} \right| \log(d_1)$$

(Chickering et al '10)

Coulomb blockade

- Droplet capacity stops the electron
- Bias & $T \sim 0$: needs energy matching

$$E(n+1, S) = E(n, S)$$

➔ current peak

- energy deformation by $\Delta S \sim \Delta Q_{\text{bkg}}$

$$E(n, S) = \frac{v}{R} \frac{(\lambda + pn - \sigma)^2}{2p} \propto (Q - Q_{\text{bkg}})^2$$



$$\Delta\sigma = \frac{B\Delta S}{\Phi_0} = \frac{1}{\nu}, \quad \Delta S = \frac{e}{n_0}$$

$U(1)$ ➔ equidistant peaks

$U(1) \times \frac{G}{H}$ ➔ modulated pattern

$$\Delta\sigma_m^\ell = \frac{1}{\nu} + \frac{v_n}{v} (h_{m+2}^\ell - 2h_m^\ell + h_{m-2}^\ell)$$

$$\frac{v_n}{v} \sim \frac{1}{10}$$



- compares states in the same sector
- spectroscopy of lowest CFT states
- $T = 0$: cannot distinguish NA state from "parent" Abelian state

$$\theta_a^\ell = \sum_{\beta=0}^{k-1} K_{a+\beta(k+2)} \chi_{a+2\beta}^\ell$$

(Bonderson et al. '10)

$T > 0$ corrections

$$\langle Q \rangle_T \sim \frac{\partial}{\partial V_o} \log \theta_a^\ell$$

- **two scales:** $0 < T_n < T_{ch}$, $T_n = \frac{v_n}{R}$, $T_{ch} = \frac{v}{R} \sim 10 T_n$

$$T < T_n : \quad \Delta \sigma_m^\ell = \dots + \frac{T}{T_{ch}} \log \left(\frac{(d_m^\ell)^2}{d_{m+2}^\ell d_{m-2}^\ell} \right), \quad d_m^\ell \text{ multiplicity of neutral states}$$

in (331) & Anti-Pfaff, not in Pfaff

$$T_n < T < T_{ch} : \quad \dots + \propto \frac{T}{T_{ch}} e^{-h_1^1 T/T_n} \frac{S_{l1}}{S_{l0}}, \quad S \text{ matrix of non-Abelian part}$$



test non-Abelian part of disk partition function

(Stern et al., Georgiev, AC et al. '09, '10)

$T > 0$ off-equilibrium

- energy offset $\Delta E_\sigma > 0$ and bias $\Delta V_o > 0$

relevant regime: $T < \Delta V_o < \Delta E_\sigma$

 thermal-activated Coulomb-Blockade conduction

$$\Gamma \sim d_a^\ell (\Delta E_\sigma^2 - \Delta V_o^2) e^{-\beta(\Delta E_\sigma - \Delta V_o)}$$

- real-time experiment of peak counting
- sensible to level multiplicity d_a^ℓ (but qualitative)

Conclusions

- non-Abelian anyons could be seen
- partition function:
 - it is simple enough
 - it defines the CFT, its sectors, fusion rules etc.
 - it is useful to compute observables
 - it can be the basis for further model building